



# STATE OF CONNECTICUT

DEPARTMENT OF PUBLIC HEALTH

## IMMUNIZATION REQUIREMENTS FOR ENROLLED STUDENTS IN CONNECTICUT SCHOOLS 2016-2017 SCHOOL YEAR



### PRESCHOOL

|               |   |
|---------------|---|
| DTaP:         | 4 doses (by 18 months for programs with children 18 months of age)  |
| Polio:        | 3 doses (by 18 months for programs with children 18 months of age)  |
| MMR:          | 1 dose on or after 1 <sup>st</sup> birthday   |
| Hep B:        | 3 doses, last one on or after 24 weeks of age   |
| Varicella:    | 1 dose on or after 1 <sup>st</sup> birthday or verification of disease  |
| Hib:          | 1 dose on or after 1 <sup>st</sup> birthday   |
| Pneumococcal: | 1 dose on or after 1 <sup>st</sup> birthday   |
| Influenza:    | 1 dose administered each year between August 1 <sup>st</sup> -December 31 <sup>st</sup> (2 doses separated by at least 28 days required for those receiving flu for the first time) |
| Hepatitis A:  | 2 doses given six calendar months apart, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday  |

### KINDERGARTEN

|               |  |
|---------------|--|
| DTaP:         | At least 4 doses. The last dose must be given on or after 4 <sup>th</sup> birthday   |
| Polio:        | At least 3 doses. The last dose must be given on or after 4 <sup>th</sup> birthday   |
| MMR:          | 2 doses separated by at least 28 days, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday                             |
| Hep B:        | 3 doses, last dose on or after 24 weeks of age   |
| Varicella:    | 2 doses separated by at least 3 months-1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday; or verification of disease |
| Hib:          | 1 dose on or after 1 <sup>st</sup> birthday for children less than 5 years old   |
| Pneumococcal: | 1 dose on or after 1 <sup>st</sup> birthday for children less than 5 years old   |
| Hepatitis A:  | 2 doses given six calendar months apart, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday                           |

### GRADES 1-4

|              |   |
|--------------|---|
| DTaP/Td:     | At least 4 doses. The last dose must be given on or after 4 <sup>th</sup> birthday. Students who start the series at age 7 or older only need a total of 3 doses. |
| Polio:       | At least 3 doses. The last dose must be given on or after 4 <sup>th</sup> birthday  |
| MMR:         | 2 doses separated by at least 28 days, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday  |
| Hep B:       | 3 doses, last dose on or after 24 weeks of age  |
| Varicella:   | 2 doses separated by at least 3 months-1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday; or verification of disease                                      |
| Hepatitis A: | 2 doses given six calendar months apart, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday  |

### GRADE 5

|            |   |
|------------|---|
| DTaP/Td:   | At least 4 doses. The last dose must be given on or after 4 <sup>th</sup> birthday. Students who start the series at age 7 or older only need a total of 3 doses. |
| Polio:     | At least 3 doses. The last dose must be given on or after 4 <sup>th</sup> birthday  |
| MMR:       | 2 doses separated by at least 28 days, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday  |
| Hep B:     | 3 doses, last dose on or after 24 weeks   |
| Varicella: | 2 doses separated by at least 3 months-1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday; or verification of disease                                      |

## GRADE 6

|            |   |
|------------|---|
| DTaP/Td:   | At least 4 doses. The last dose must be given on or after 4 <sup>th</sup> birthday. Students who start the series at age 7 or older only need a total of 3 doses. |
| Polio:     | At least 3 doses. The last dose must be given on or after 4 <sup>th</sup> birthday  |
| MMR:       | 2 doses separated by at least 28 days, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday  |
| Hep B:     | 3 doses, last dose on or after 24 weeks of age  |
| Varicella: | 1 dose on or after 1 <sup>st</sup> birthday; or verification of disease   |

## GRADES 7-12

|                |  |
|----------------|--|
| Tdap/Td:       | 1 dose for students who have completed their primary DTaP series. Students who start the series at age 7 or older only need a total of 3 doses of tetanus-diphtheria containing vaccine, one of which must be Tdap |
| Polio:         | At least 3 doses. The last dose must be given on or after 4 <sup>th</sup> birthday   |
| MMR:           | 2 doses separated by at least 28 days, 1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday   |
| Meningococcal: | 1 dose   |
| Hep B:         | 3 doses, last dose on or after 24 weeks of age   |
| Varicella:     | 2 doses separated by at least 3 months-1 <sup>st</sup> dose on or after 1 <sup>st</sup> birthday; or verification of disease   |

- DTaP vaccine is not given on or after the 7<sup>th</sup> birthday and may be given for all doses in the primary series.
- Tdap can be given in lieu of Td vaccine for children 7 years and older unless contraindicated. Tdap is only licensed for one dose.
- Hib is not required for children 5 years of age or older.
- Pneumococcal is required for all Pre-K and K students less than 5 years of age.
- Hep A requirement for school year 2016-17 applies to all Pre-K, K, 1<sup>st</sup>, 2<sup>nd</sup>, 3<sup>rd</sup> & 4<sup>th</sup> graders born 1/1/07 or later.
- Hep B requirement for school year 2016-2017 applies to all students in grades K-12.  
Spacing intervals for a valid Hep B series: at least 4 weeks between doses 1 and 2; 8 weeks between doses 2 and 3; at least 16 weeks between doses 1 and 3; dose 3 should not be given before 24 weeks of age.
- Second MMR for school year 2016-2017 applies to all students in grades K-12.
- Meningococcal Conjugate requirement for school year 2016-17 applies to all students in grades 7-12
- Tdap requirement for school year 2016-17 applies to all students in grades 7-12
- If two live virus vaccines (MMR, Varicella, MMRV, Intra-nasal Influenza) are not administered on the same day, they must be separated by at least 28 days (there is no 4 day grace period for live virus vaccines). If they are not separated by at least 28 days, the vaccine administered second must be repeated.
- Lab confirmation of immunity is **only** acceptable for Hep A, Hep B, Measles, Mumps, Rubella, and Varicella.
- **VERIFICATION OF VARICELLA DISEASE:** Confirmation in writing by a MD, PA, or APRN that the child has a previous history of disease, based on family or medical history.
- For the full legal requirements for school entry visit [www.ct.gov/dph/cwp/view.asp?a=3136&Q=467374&PM=1](http://www.ct.gov/dph/cwp/view.asp?a=3136&Q=467374&PM=1)
- If you are unsure if a child is in compliance, please call the Immunization Program at (860) 509-7929.

### **New Entrant Definition:**

\*New entrants are any students who are new to the school district, including **all** preschoolers and all students coming in from Connecticut private, parochial and charter schools located in the same or another community. **All pre-schoolers, as well as all students entering kindergarten**, including those repeating kindergarten, and those moving from any public or private pre-school program, even in the same school district, **are considered new entrants**. The one exception is students returning from private approved special education placements—they are not considered new entrants.

### **Commonly Administered Vaccines:**

| <u>Vaccine:</u> | <u>Brand Name:</u> | <u>Vaccine:</u> | <u>Brand Name:</u>                            |
|-----------------|--------------------|-----------------|---|
| DTaP-IPV-Hib    | Pentacel           | MMRV            | ProQuad                                       |
| DTaP-HIB        | TriHibit           | PCV7            | Prennar                                       |
| HIB-Hep B       | Comvax             | PCV13           | Prennar 13                                    |
| DTaP-IPV-Hep B  | Pediarix           | DTaP-IPV        | Kinrix  |
| Hepatitis A     | Havrix, Vaqta      | Influenza       | Fluzone, FluMist, Fluviron, Fluarix, FluLaval |